

APPENDIX C

Standard Operating Procedures for Fertility Control Treatment

The following management and monitoring requirements are part of the Proposed Action:

- PZP vaccine would be administered by trained BLM personnel.
- The fertility control drug is administered with two separate injections: (1) a liquid dose of PZP is administered using an 18 gauge needle primarily by hand injection; (2) the pellets are preloaded into a 14 gauge needle. These are loaded on the end of a trocar (dry syringe with a metal rod) which is loaded into the jabstick which then pushes the pellets into the breeding mares being returned to the range. The pellets and liquid are designed to release the PZP over time similar to a time release cold capsule.
- Delivery of the vaccine would be as an intramuscular injection while the mares are restrained in a working chute. 0.5 cubic centimeters (cc) of the PZP vaccine would be emulsified with 0.5 cc of adjuvant (a compound that stimulates antibody production) and loaded into the delivery system. The pellets would be loaded into the jabstick for the second injection. With each injection, the liquid and pellets would be propelled into the left hind quarters of the mare, just below the imaginary line that connects the point of the hip and the point of the buttocks.
- All treated mares would be freeze-marked on the hip to enable researchers to positively identify the animals during the research project as part of the data collection phase.
- At a minimum, monitoring of reproductive rates using helicopter flyovers will be conducted in years 2 through 4 by checking for presence/absence of foals. The flight scheduled for year 4 will also assist in determining the percentage of mares that have returned to fertility. In addition, field monitoring will be routinely conducted as part of other regular ground-based monitoring activities.
- A field data sheet will be used by the field applicators to record all the pertinent data relating to identification of the mare (including a photograph when possible), date of treatment, type of treatment (1 or 2 year vaccine, adjuvant used) and HMA, etc. The original form with the data sheets will be forwarded to the authorized officer at National Program Office (NPO) (Reno, Nevada). A copy of the form and data sheets and any photos taken will be maintained at the field office.
- A tracking system will be maintained by NPO detailing the quantity of PZP issued, the quantity used, disposition of any unused PZP, the number of treated mares by HMA, field office, and state along with the freeze-mark applied by HMA.
- The field office will assure that treated mares do not enter the adoption market for three years following treatment. In the rare instance, due to unforeseen circumstance, treated mare(s) are removed from an HMA before three years has lapsed, they will be maintained in either a BLM facility or a BLM-contracted long term holding facility until expiration of the three year holding period. In the event it is necessary to remove treated mares, their removal and disposition will be coordinated through NPO. After expiration of the three year holding period, the animal may be placed in the adoption program or sent to a long-term holding facility.